	3			27 Recid PCI/PIC 07 NOV 2000				
FOR (RE)	M PTO / 11-98)	-1390 US DEPARTMENT C	F COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER 1303-110				
	TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) U.S. APPLICATION NO. (If Known, see 37 C F R 1.5)							
	CONCERNING A FILING UNDER 35 U.S.C. 371 U 7/(TOBA Als @n@i)							
INT	ERNAT	FIONAL APPLICATION NO.	INTERNATIONAL FILING DATE	PRIORITY DATE CLAIMED				
-	PCT/EP99/03075 5 May 1999 8 May 1998							
TITI	TITLE OF INVENTION GENETICALLY-MODIFIED FIBROBLASTS AND THE USE THEREOF							
APF	APPLICANT(S) FOR DO/EO/US							
<u> </u>	MAVILIO							
App	Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:							
1.	\boxtimes	This is a FIRST submission	of items concerning a filing under 35 U.S.C. 3	71.				
2.		This is a SECOND or SUBS	SEQUENT submission of items concerning a fil	ing under 35 U.S.C. 371.				
3.	\boxtimes		o begin national examination procedures (35 L tion of the applicable time limit set in 35 U.S.C.					
4.	The state of the s							
5.	5. A copy of the International Application as filed (35 U.S.C. 371(c)(2)).							
14								
6	<u> </u>							
	7.0 Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).							
14	i are transmitted herewith (required only if not transmitted by the International Bureau). b							
8.1		A translation of the amendm	nents to the claims under PCT Article 19 (U.S.C	C. 371(c)(3)).				
9.	\boxtimes	An oath or declaration of the	e inventor(s) (35 U.S.C. 371(c)(4)).					
	10/4 A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).							
Iten	Items 11. To 16. Below concern document(s) or information included:							
11.		An Information Disclosure S	tatement under 37 C.F.R. 1.97 and 1.98.					
12.	×		An assignment document for recording. A separate cover sheet in compliance with 37 C.F.R. 3.28 and 3.31 is included.					
13.	\boxtimes	A FIRST preliminary amend A SECOND or SUBSEQUE						
14.		A substitute specification.						
15.		A change of power of attorn	ey and/or address letter.					
	Other items or information. Verified Statement Claiming Small Entity Status - Independent Inventor & Small business Concern/ PTO-1449/ International Search Report							

\$26 Rec'd PCT/PTO 07 NOV 2000

							3 UTNO		
U.S. APPLICATION NOW (If known),	#855315	İ	PCT/EP99/03075			TTC	1303-110	NUM	BER
17. The following fees						C/	ALCULATIONS	PTO	USE ONLY
BASIC NATIONAL FEE Neither international nor international sea	(37 C.F.R. 1.4 preliminary exa rch fee (37 C.F	minatio .R. 1.44)-(5): in fee (37 C.F.R. 1.482) i5(a)(2)) paid to USPTO ed by the EPO or JPO		1000 00				
International prelimin	ary examinatio	n fee (3	7 C.F.R. 1.482) not paid to)					
International prelimin	ary examinatio	n fee (3	7 C.F.R. 1.482) not paid to	USPTO					
International prelimin	ary examinatio	n fee pa	5(a)(2) paid to USPTO aid to USPTO (37 C.F.R. 1	.482)					
International prelimin	ary examinatio	n fee pa	CT Article 33(1)-(4) aid to USPTO (37 C.F.R. 1	.482)					
and all claims satisfic	ed provisions of	PCT A	rticle 33(1)-(4)			s	860.00	г	
Surcharge of \$130.00 for fu	rnishing the oat	h or de		□ 30	AWOUNT =	3	860.00	<u> </u>	
months from the earliest cla		ate (37 (ATE	\$	0.00	L	
Total Claims	10	-20 =	0	X	\$18.00	s	0.00	_	
Independent Claims	1	-3 =	0	X	\$80.00	Ť	0.00	-	
MULTIPLE DEPENDENT C					0.00	s	0.00		
			TOTAL OF AB			\$	860.00		
Reduction by ½ for filing by must also be filed (Note 37			le. A Small Entity Statem				430,00		
4.3	<u> </u>	,		s	UBTOTAL =	\$	430.00		
Processing fee of \$130.00,	for furnishing th	e Engli	sh Translation later than	20 🗌 30					
months from the earliest cla	imed priority da	ate (37 (+		L	0.00	_	
N					ONAL FEE =	\$	430.00		
Fee for recording the enclose							40.00		
accompanied by an approp	late cover snee	bonder	.F.H. 3.28, 3.31). \$40.00	Small Entir	+ + + + + + + + + + + + + + + + + + + +	\$	0.00	├~	
Let for Petition to Nevive U	nintentionally A	Danuoi	TOT	AI FEES E	NCLOSED =	\$	470 .00	┝	
tu				ALILLOL	HOLOGED =		mount to be:		
a						l ^`	refunded	l s	
(a b						Г	Charged	\$	
Please charge m form is enclosed. The Commission Deposit Account The entire conten application. NOTE: Where an appropr	r Deposit Accor er is hereby aut No. <u>14-1140</u> . A t of the foreign iate time limit nust be filed an	unt No. horized duplica applica under 3	ver the above fees is enclot- 14-1140 in the amount of 3 to charge any additional finate copy of this form is encition(s), referred to in this a 37 C.F.R. 1.494 or 1.495 head to restore the application of the application o	to co ees which m losed. pplication is	ay be required are hereby ind met, a petiti ding status.	l, or	credit any ove	rpay	ment to
NIXON & VANDERHYE P.O).			SIGNATE	/ L		1		
1100 North Glebe Road, 8th									
Arlington, Virginia 22201									
Telephone: (703) 816-4000				NAME	R. Crawford	_			
}				05.05-			M	-	
				25,327	RATION NUMBE	-0	November Date	7, 20	טטנ
				neulsti	TATION NUMBE	<u>- ri</u>	Date		

Q	*	S	1	V	

			•	
				OMB No. 0651-0011 (12/31/86)
	ant or Patentee:	-		Attorney's. Dkt. No.
	or Patent No.:			
For:	or Issued:			
101.				
		STATU	TATEMENT (DECLARATION) CLA S [37 19(f) and 1.27(c)] - INDEPEN	DENT INVENTOR
paying the inv	reduced fees ur ention entitled:	nder Section 41 (a) a	and (b) of Title 35, United States Co	inventor as defined in 37 CFR 1.9(c) for purposes of de, to the Patent and Trademark Office with regard to
		ibroblasts and the	use thereof	
descril	oed in			
	the specification	n filed herewith.		
	application Seri	al No.	, filed	
	patent No.		, issued	
license person	e, any rights in th i had made the i	e invention to any pe	erson who could not be classified as	on under contract or law to assign, grant, convey or an independent inventor under 37 CFR 1.9(c) if that a small business concern under 37 CFR 1.9(d) or a
Each p	ct or law to assign	n, grant, convey, or li	cense any rights in the invention is li	veyed, or licensed or am under an obligation under sted below:
(N 4.1		erson, concern, or or oncerns or organizat		
NOTE:	Separate verifie g to their status a	d statements are re as small entities. (37	quired from each named person, c CFR 1.27)	oncern or organization having rights to the invention
LÚ	Name			
4	Address			
44		☐ Individual	☐ Small Business Concern	☐ Nonprofit Organization
pris.	Name			
O	Address			
44		☐ Individual	☐ Small Business Concern	☐ Nonprofit Organization
()	Name			
()	Address			The state of the s
	Addicas	Individual	☐ Small Business Concern	☐ Nonprofit Organization
			_ small pasitions contour	
entity s	tatus prior to pay	ring, or at the time of	ation of patent, notification of any ch f paying, the earliest of the issue fee ate. [37 CFR 1.28(b)]	ange in status resulting in loss of entitlement to small or any maintenance fee due after the date on which
made a statem	ieved to be true; are punishable by ents may jeopard	and further that thes fine or imprisonme	se statements were made with the k nt, or both, under § 1001 of Title 18	and that all statements made on information and belief nowledge that willful false statements and the like so of the United States Code, and that such willful false reon, or any patent to which this verified statement is
directe	d.		. "	
	MAVILIO			The second secon
	NAME OF I	VENIOR OF	NAME OF INVENTOR	NAME OF INVENTOR
	Signal	ture	Signature	Signature
13	3-10-2000		- Oignature	Signature
	DAT		DATE	DATE

OMB No. 0651-0011 (12/31/86)

Applicant or Patentee	Attomey's.Dkt. No.
Serial or Patent No.:	
Filed or Issued:	
For:	VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY
	STATUS [37 19(f) and 1.27(c)] - SMALL BUSINESS CONCERN
I hereby declare that I	
□ the o	er of the small business concern identified below:
	al of the small business concern empowered to act on behalf of the concern identified below:
	OF CONCERN GENERA S.p.A.
ADDI	SS OF CONCERN Via Olgettina, 58
I because the street and the st	MILANO, Italy
CFR 121.3-18, and re 35, United States Coo 500 persons. For pur previous fiscal year of pay periods of the fisc	a above identified small business concern qualifies as a small business concern as defined in 13 oduced in 37 CFR 1.9(d), for purposes of paying reduced fees under section 41(a) and (b) of Title in that the number of employees of the concern, including those of its affiliates, does not exceed ses of this statement (1) the number of employees of the business concern is the average over the le concern of the persons employed on a full-time, part-time or temporary basis during each of the year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern or to control the other, or a third party or parties controls or has the power to control both.
I hereby declare that identified above with n	ghts under contract or law have been conveyed to and remain with the small business concern and to the invention, entitled:
Genetically-modified	broblasts and the use thereof
by inventors	de-called in
	filed herewith. described in
	No , filed
patent No.	, issued
having rights to the in- who could not qualify concern which would EFR 1.9(e). *NOTE: S rights to the invention Name	above-identified small business concern are not exclusive, each individual, concern or organization into is listed below* and no rights to the invention are neld by any person, other than the inventor is an independent inventor under 37 CFR 1.9(c) if that person made the invention, or by any it qualify as a small business concern under 37 CFR 1.9(d), or a nonprofit organization under 37 parate verified statements are required from each named person, concern or organization having erring to their status as small entities (37 CFR 1.27)
Address	
£.2	Individual Small Business Concern Nonprofit Organization
Name	
Address	
	Individual Small Business Concern Nonprofit Organization
entitlement to small er fee due after the date I hereby declare that information and belief false statements and	to file, in this application of patent, notification of any change in status resulting in loss of by status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance which status as a small entity is no longer appropriate. [37 CFR 1.28(b)] ill statements made herein of my own knowledge are true and that all statements made on e believed to be true; and further that these statements were made with the knowledge that willful like so made are punishable by fine or imprisonment, or both, under § 1001 of Title 18 of the
thereon, or any patent	that such willful false statements may jeopardize the validity of the application, any patent issuing which this verified statement is directed.
NAME OF PERSON S TITLE OF PERSON O	
ADDRESS OF PERSO	
SIGNATURE	DATE 13.10.2000

09/674853 526 Rec'd PCT/PTO 07NOV 2888

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of

MAVILIO

Atty. Ref.:

1303-110

Serial No.

(To Be Assigned)

Group:

National Phase of F

PCT/EP99/03075

Filed:

November 7, 2000

Examiner:

For: GENETICALLY-MODIFIED FIBROBLASTS AND THE

USE THEREOF

Examiner.

November 7, 2000

Assistant Commissioner for Patents Washington, DC 20231

Sir:

PRELIMINARY AMENDMENT

Prior to calculation of the filing fee and in order to place the above identified

application in better condition for examination, please amend the claims as follows:

IN THE CLAIMS

Claim 9, lines 1-2, change "claims 1-8" to --claim 1--.

REMARKS

The above amendments are made to place the claims in a more traditional format.

Respectfully submitted,
NIXON & VANDERHYE P.O.

By:

Arthur R/Crawford Reg. No. 25,327

ARC:ms

1100 North Glebe Road, 8th Floor Arlington, VA 22201-4714 Telephone: (703) 816-4000

Facsimile: (703) 816-4100

10

15

20

7/07/853

PTO/PCT Rec'd 7 NOV 2001

GENETICALLY-MODIFIED FIBROBLASTS AND THE USE

THEREOF

The present invention relates to an ex-vivo method for the myogenic conversion of genetically-modified fibroblasts, for use in the correction of congenital defects of the muscular system, such as primary monogenic myopathies, or in the production of therapeutical proteins. Furthermore, the invention relates to the use of genetically-modified fibroblasts for the preparation of compositions for cell implants for the treatment of muscle pathologies with a genetic aetiology, or for the in vivo secretion of recombinant proteins of therapeutical importance. In particular, the invention is valuable for the treatment of muscular dystrophies, a heterogeneous group of severe degenerative diseases of the muscle caused by mutations in the genes encoding for the membrane-associated protein dystrophin or for other members of the dystrophin-associated protein complex connecting the muscle fiber cytoskeleton with the extracellular matrix.

Until now, the attempts to correct muscular dystrophy with gene therapy based on the implantation of muscles or other tissues stem cells, have been thwarted by enormous difficulties in the isolation of a suitable number of myogenic cells to be genetically-modified or to the poor yield of phenotypic conversion. By way of example, the difficulty in obtaining a reasonable number of satellite cells to be genetically-modified is a limit to gene therapy of patients with Duchenne muscular dystrophy or with Becker muscular dystrophy (Salvatori, G., et al., 1993; Webster, C. et al., 1990). Further problems related to the implantation of genetically-modified cells in dystrophic patients are the immunogenicity of viral

15

20

25

WO 99/58646 PCT/EP99/03075

vectors, the difficulty in developing an effective administration route and the poor survival of injected cells. Furthermore, the expression of dystrophin itself has been reported to induce cellular and/or humoral immune responses in dystrophin-lacking patients (New England J. of Medicine 333, 732-733, 1995).

It has been known for some time that forced expression of MyoD, and related members of the myogenic family of basic helix-loop-helix transcription factors, would activate myogenesis in non-muscle cells (Davis, R. L., et al., 1987; Cossu, G., et al., 1996; Cossu, G., 1997). The Applicant itself has recently reported that fibroblasts undergo myogenic differentiation when co-cultured with myogenic lines (C2C12) or primary cells, but not with other cell types (Breton, M. et al., 1995; Gibson, A.J., et al., 1995; Salvatori, G. et al., 1995). WO 95/12979 discloses a general method for inducing cell differentiation to a novel phenotype, for example muscular phenotype, by inserting the genic sequences of factors such as MyoD, myogenin, Myf-5 and MRF4 into the cells. The same document neither considers any possibility to convert genetically-modified cells to the myogenic phenotype in order to make them suitable for gene therapy, nor envisages the possibility to carry out an "ex vivo" myogenic conversion.

GB 2293604 discloses the use of fibroblasts in the treatment of muscular pathologies, wherein fibroblasts themselves can be transformed with a therapeutically active gene, for example the dystrophin gene, whereas the possibility to induce differentiation by means of myogenic factors has not been considered.

In light of what stated above, it appears remarkably advantageous to rely on a method which allows a high number of cells to be converted

15

20

25

to the myogenic phenotype, after being genetically-modified ex-vivo, in order to correct a genetic defect, such as muscular dystrophy, or any other dysfunction related, for instance, to insufficient or incorrect production of a protein or a plasma factor or other secreted or circulating proteins, as in the case of diabetes, hemophilia, pituitary dwarfism.

It has now been found that it is possible to convert genetically-modified fibroblasts to the muscular phenotype, with a high-efficiency process, so as to effectively produce therapeutical proteins or proteins able to correct the genetic defects, particularly muscular dystrophy.

According to a first aspect, the invention provides a high-efficiency method for the preparation of genetically-modified fibroblasts expressing muscle lineage commitment genes, which comprises:

- ex-vivo transduction of fibroblasts with a therapeutic gene or a gene capable of correcting a genetic defect;
- b) transient expression of the muscle lineage commitment gene in fibroblasts transduced as at point (a), through transformation of the cells with a high efficiency DNA transfer method, preferably a viral vector, more preferably selected from baculovirus, adenoassociated viruses and, most preferably, adenovirus, or with other high efficiency DNA transfer means, such as liposomes, nucleoprotein complexes, polyethylenimines, wherein the gene of the muscle lineage commitment is under the control of a strong promoter, preferably a viral promoter.

"Muscle lineage commitment gene" means any gene capable of converting fibroblasts to the myogenic phenotype, particularly MyF-5, MRF4, myogenin and, preferably, MyoD genes.

Therapeutical genes which can be transduced into fibroblasts

14

9

10

15

20

25

WO 99/58646 PCT/EP99/03075

before the myogenic conversion are, for example, genes involved the various types of muscle dystrophies, such as sarcoglycan, emerin and, preferably, dystrophin genes, or genes coding for circulating plasma proteins, such as coagulation factors or insulin, for hormones, such as growth hormone, or other genes, such as hypoxanthine-guanine phosphoribosyl transferase, adenosine deaminase, purine nucleoside phosphorylase, glucocerebrosidase, low density lipoprotein receptor, phenylalanine hydroxylase, arginine succinate synthetase and aryl-sulfatase.

A second aspect of the invention relates to genetically-modified fibroblasts, transiently expressing a gene of the muscle lineage commitment, preferably MyoD, obtained by the method of the invention.

The conversion of fibroblasts to the myogenic phenotype by transient expression of a muscle lineage commitment gene, proved to be surprisingly more effective than other alternative techniques. In fact, coculture of fibroblasts with satellite cells, treatment with dexamethasone transfection with calcium phospnate, lipofectamine. electroporation of the same plasmid expressing MyoD, induced myogenic conversion in a percentage ranging from 1% to 14%, in comparison with values higher than 70% in case of infection with a recombinant adenoviral vector. Furthermore, the efficiency of conversion was found to increase linearly for m.o.i. ("multiplicity of infection") values from 500 to 2000, whereas cytotoxicity at maximum m.o.i. values, remained within acceptable limits. Similar results were obtained with human or murine fibroblasts, whereas as far as tissutal fibroblasts are concerned, dermal fibroblasts gave better results. One main advantage of the method of the invention is that transient expression of the exogenous muscle lineagecommitment gene activates the corresponding endogenous gene, therefore

15

20

irreversibly committing cells to myogenesis, thus rendering constitutive expression of the transgene unnecessary. The actual myogenic conversion of fibroblasts transiently expressing exogenous MyoD has been confirmed by immuno-histochemistry, electron microscopy and gene expression analysis of different muscle-specific markers, such as myosin light and heavy chains, acetylcholine receptor α -subunit, M creatine kinase, myogenin and MyoD.

The high efficiency of the method allows the preparation of genetically manipulated fibroblasts in which a therapeutic gene, or a gene correcting the genetic defect in the muscle has been inserted, which may be stably converted to the myogenic phenotype. In fact, conversion to the muscle phenotype prevents cell replication and provides stable implants, based on the use of muscle fibers.

Furthermore, the efficiency of the method as well as the stability of the resulting product allow to successfully carry out gene-therapy, in which genetically-modified fibroblasts expressing a gene of the muscle lineage commitment, after appropriate ex-vivo treatment, are injected into muscle tissue where they are capable of regenerating the muscle fibers and express the correct gene.

A number of evidences proved that fibroblasts treated ex-vivo as herein disclosed are able to regenerate muscle fibers in mice, indistinguishable from those originating from primary myogenic (satellite) cells, with a long-lasting effect. Moreover, after treatment, only antibody mediated immune response against proteins of adenoviral infected fibroblasts has been detected. No cell-mediated immune response has been observed at the site of injection of the modified fibroblasts. This is a significant advantage in gene therapy, as one of the major problems

associated to the in vivo use of adenoviral vectors is the induction of significant, T-cell-mediated immune responses against both the viral and transgenic proteins.

According to a further aspect, the invention relates to the use of 5 fibroblasts obtained by the above described method, for "ex vivo" gene therapy or for the preparation of compositions for stable cell implants based on muscle fibers.

A typical application of the ex-vivo gene therapy comprises, for example, isolating dermal fibroblasts from a patient suffering from a genetic disease which alters the muscle structure or functionality, such as dystrophy, expanding the cultured cells, transducing the cells ex-vivo with a retroviral vector containing the correct gene, and inducing myogenesis by infection with an adenoviral vector, or with another vector as mentioned above, containing the gene of the muscle lineage commitment, and re-implanting modified cells into the muscle tissue.

10

20

10

Another typical approach for the preparation of cell implants based on muscle fibers comprises isolating dermal fibroblasts from a patient suffering from a disease characterised by, for example, lack of a plasma protein (such as diabetes, pituitary dwarfism, hemophilia), expanding the cultured cells, transducing the cells ex-vivo with a retroviral vector containing the therapeutic gene, and inducing myogenesis by infection with an adenoviral vector, or with another vector as mentioned above, containing the muscle lineage commitment gene, encapsulating the modified cells in the most suitable implantation matrix and implanting the preparation in the preferred body site. In this specific application, induction of myogenesis and subsequent differentiation into muscle tissue is not aimed at the restoration of muscle functionality, but to the stability

of the implant by arrest of cell division.

As further applications of the present invention, its use in zootechnic field can be envisaged, to induce a specific characteristic in animals, for example an increase in weight, by production of a hormone or any other substance by the cells treated ex vivo with the herein disclosed method and then re-implanted into the animal; or the creation or use of animal models of human pathologies for the study of novel therapeutical procedures to be carried out with the method of the invention.

The following examples illustrate the invention in greater details.

Example 1

10

15

20

25

Preparation of genetically-modified fibroblasts expressing MyoD.

Fetal fibroblasts were isolated from skin and skeletal muscle of C3H mouse embryos (15-17 days) or legally aborted human fetuses (8-12 weeks) as described in Salvatori et al., *Human Gene Ther.*, 1993, 4:713-723, or human dermal or muscle fibroblasts were obtained from tissues of patients undergoing post-traumatic surgery. In this case tissues were finely minced with scissors, digested with 2 mg/ml dispase, 0.1 mg/ml collagenase in phosphate buffered saline (PBS) for 45' at 37°C, washed in RPMI medium, and pipetted to obtain a single-cell suspension.

Murine bone marrow fibroblasts transgenic for the lacZ gene with nuclear localization, were obtained by resuspending and plating cells flushed from the long bones of 8-10-wk-old MLC3F/nlacZ mice. Human bone marrow fibroblasts were obtained from healthy donors after removal of non-adherent cells. All cells were grown in RPMI supplemented with 15% Fetal calf serum (FCS), 1% gentamycin and 0.3 mM ß-mercaptoethanol (growth medium). Human bone marrow fibroblasts were

1

14

10

10

20

supplemented with 2 ng/ml bFGF.

Myogenic differentiation was induced by shifting the cells to RPMI supplemented with 2% horse serum (differentiation medium). Fibroblasts were purified by sub-culture (at least two rounds) in growth medium.

5 Removal of myogenic cells was routinely verified by immunocytochemical staining of a cell aliquot sub-cultured for 5 days in differentiation medium.

For transduction, fibroblasts (from third to tenth passage) were infected with a replication-defective retroviral vector (LBSN) expressing a cytoplasmic β-galactosidase gene under the LTR promoter as described (Salvatori, G., et al., 1993).

For the subsequent step (transient expression of MyoD), cells were treated with the Ad5-derived, E1A-deleted adenoviral vector expressing the full-length murine MyoD c-DNA under the transcriptional control of Rous Sarcoma Virus (RSV) LTR (Murry, C. E., et al., 1996). In some control experiments, cells were transfected by standard precipitation techniques with calcium-phosphate or lipofectamine (Dotap) with 10 µg of the PMC11 plasmid, containing the MyoD cDNA under the control of the CMV promoter.

Alternatively, cells were electroporated with 6 μg of the same plasmid in growth medium at 120 V, 960 mF. After transduction, cells were grown for 24 hrs in growth medium and then either shifted to differentiation medium for 3-4 days or injected in vivo.

In some experiments, cells were pre-labelled with 0.5 mCi/ml of [14C] thymidine (Amersham) for 24 h and then exposed to the adenoviral vector expressing MyoD and survival was measured by counting residual cpm incorporated.

15

25

Example 2

Implantation of MyoD-converted fibroblasts into mouse muscles.

1 or 2 millions of transduced (or transgenic) human or murine fibroblasts, converted by transient expression of MyoD, were trypsinized, 5 resuspended in 20-50 μl of PBS and injected into a single site of the regenerating tibialis anterior muscle of either syngeneic (C3H) or immunodeficient (scid/bg) mice which had received a 30-μl injection of 10-5 M cardiotoxin (Latoxan) 48 hrs earlier.

For some experiments, myogenically converted mouse fibroblasts were labelled with DiI dye 1 hr before injection in vivo (a 0.5% solution of DiI in absolute ethanol was diluted just before use in 0.3 M sucrose to a final concentration of 0.05%).

Mice were sacrificed after various periods of time, and muscles were cryo-sectioned and stained for β-galactosidase activity or processed for immuno-hystochemistry. At the time of sacrifice, serum was collected from C3H mice and reacted with adeno-infected or control cells at various dilutions, followed by a fluorescein-conjugated anti-mouse IgG secondary antibody.

Example 3

20 Immunocytochemistry, electron microscopy and RNA analysis techniques.

Immunofluorescence analysis was carried out as described (Cusella - De Angelis, M.G., et al., 1994) using the following antibodies: MF20, a monoclonal antibody which recognises all sarcomeric myosins (Bader, D., et al., 1982); a rabbit antiserum against sarcomeric proteins (Tajbakhrsh, S., et al., 1994); an anti-MyoD polyclonal antibody (Hasty, P., et al., 1993); BD5, a monoclonal antibody which recognises slow myosin heavy

20

chains (Yasin, R., et al., 1977); a rabbit polyclonal antibody against human fetal myosin (Edom, F., et al., 1994); an anti-bromo-deoxyuridine (BrdU) monoclonal antibody; anti-leu, anti-CD4 and anti-Mac3 rat anti-mouse leukocyte antigen antibodies. Briefly, cells cultures or cryostat 5 - sections were fixed with 4% paraformaldehyde for 10 min at 4°C, washed 3X in PBS, and incubated at 4°C with the primary antibodies in PBS + 1% bovine serum albumin (BSA). BrdU labelled cell-cultures were treated for 10 min at rt with 2M HCl and washed 3X with PBS before incubation with the anti-BrdU antibody.

After the first incubation, cells or sections were washed 3X with PBS + 1% BSA and incubated for 1 hr at rt with rhodamine-conjugated goat anti-mouse Ig or with a fluorescein-conjugated goat anti rabbit Ig (1:100 dilution).

Cultures or sections were then washed, mounted in 75% glycerol/PBS (pH 7.5), and observed under a Zeiss Axiophot epifluorescence microscope.

For electron microscopy cells were washed in PBS, fixed in 2% glutaraldehyde in 0.1 M Millonig's buffer (pH 7.4) for 1 hr at 4°C, post-fixed for 1 hr in 1% buffered osmium tetroxide and dehydrated in graded alçohol. Cells were treated with propylene oxide, embedded in Epon 812 and cut into ultrathin sections. The ultrathin sections were stained with uranyl acetate and lead citrate, and examined with a Zeiss 109 electron microscope.

For RNA analysis, RNA was extracted by the method of Chomczynski et al. (Chomczynski, P., et al., 1987), run in 15 µg aliquots on 1% agarose/formaldehyde gels, and transferred by capillary Northern blotting to nylon membranes (Amersham Hybond-N).

WO 99/58646

20

25

Filters were cross-linked for 2 hrs at 80 °C under vacuum, and hybridized to [32P]-labelled probes for MyoD and myogenin (Bober, E., et al., 1991), MLC1F and MCK (Lyons, G. E., et al., 1991), Ach-receptor α-subunit (25) under standard conditions (26).

5 - Example 4

Analysis of in vitro converted fibroblasts expressing the myogenic phenotype.

In order to compare the adeno-MyoD-induced phenotype with that of primary myogenic cells, a number of muscle-specific markers were analysed by immuno-cytochemistry, electron microscopy and gene expression analysis, in human and murine converted dermal fibroblasts, and in differentiating satellite cells of comparable age.

All differentiated myotubes derived from fusion of converted murine newborn dermal fibroblasts express fast embryonic myosin heavy chains, while a fraction of them also express slow myosin heavy chains, while a fraction of them also express slow myosin heavy chains, as observed in myotubes derived from satellite cells.

Similarly, myotubes derived from human fetal dermal fibroblasts show a well organized initial sarcomerogenesis, with aligned sarcomeres and patterned Z lines, as reported for myotubes derived from primary myogenic cells, which never complete sarcomerogenesis in vitro.

In order to verify whether the introduction of MyoD in primary fibroblasts would also activate transcription of genes responsible for membrane and metabolic muscle-specific functions, the Applicant measured by Northern blotting the expression of RNAs coding muscle-specific proteins such as the acetylcholine (ACh) receptor α-subunit, M creatine kinase (MCK), as well as MyoD, myogenin, and myosin light

15

20

25

WO 99/58646 PCT/EP99/03075

chain 1 fast (MLC1F) chosen as positive controls.

No detectable level of any of these messages was detected in unconverted fibroblasts; on the other hand, these mRNAs were expressed at comparable levels in myotubes derived from converted fibroblasts and 5. in myotubes derived from satellite cells of corresponding age.

Thus, under all the parameters analysed in vitro, myotubes originating from converted fibroblasts cannot be distinguished from those derived by primary myogenic cells.

Afterwards, the Applicant investigated whether fibroblasts exposed to the MyoD adenoviral vector would still maintain the capacity to divide, since MyoD overexpression has an anti-proliferative effect. Murine and human fetal skin fibroblasts were infected with the MyoD adenoviral vector for 3 hrs at an m.o.i. of 2,000 in serum-free medium, cultured for 12 hrs in growth medium containing 10 mM BrdU, fixed at 12, 24, 48 and 72 hrs after infection, and double-stained with anti-MyoD and anti-BrdU antibodies.

24 hrs after exposure to the adenoviral vector, only a few cells expressing MyoD had incorporated BrdU, while no double-positive cells were detected at 48 and 72 hrs. Expression of MyoD therefore blocks cell division in converted fibroblasts.

Example 5

Myogenic conversion efficiency of human and murine fibroblasts of different origins

Fibroblasts were isolated, expanded, infected with the MyoD adenoviral vector for 3 hrs at an m.o.i. of 2,000, cultured for a further 24 hrs and then induced to differentiation as described. Table 1 shows the conversion % values.

5 -

10

25

The conversion efficiency was measured as percentage of cells which expressed sarcomeric myosin heavy chains, as shown by staining with anti-myosin antibody. (MF20). Values shown are the average of two separate experiments, each performed in triplicate.

Table 1

	M	urine	Hur	nan
	Fetal	Adult	Fetal	Adult
Derma	70	56	65	42
Muscle	43	44	40	32
Bone marrow	ND	6	ND	6

ND: not done.

15 Example 6

Survival of MyoD-converted fibroblasts and satellite cells (from MLC3F/LacZ mice) after injection into regenerating Tibialis Anterior muscle (TA) of SCID/bg mice.

 $20~\mu l$ containing $1x10^6$ cells were injected into a single site of 20 regenerating TA of SCID/bg mice.

At the indicated times, mice were sacrificed and successive 15 μm cryostat sections were prepared from the TA muscle. 1 every 5 slides was X-gal stained. Nuclei were counter-stained with Hoechst. The X-gal stained sections, were scored for the number of β -gal+ nuclei and the obtained values were multiplied x 5 (total No. of nuclei). In addition β -gal+/total nuclei in 200 μ^2 areas was counted. Results are reported in Table 2. Note that the increase in the β -gal+/total nuclei ratio is due to the

1)

decrease in monunucleated cells which progressively occurs during muscle regeneration (fusion).

Table 2

	No. of B-gal ⁺ nuclei/muscle (0% of injected cells)	uclei/muscle cells)	No. of B-gal^+ nuclei/injection (200 μ^2) (0% of donor nuclei)	No. of B-gal ⁺ nuclei/injection site (200 μ^2) (0% of donor nuclei)
Time(weeks)	MyoD	satellite cells	MyoD	satellite cells
	fibroblasts		fibroblasts	
0	1.0x106 (100)	1.0x106 (100)		
1	$2.1x10^3$ (0.2)	$0.5 \times 10^3 (0.05)$	8/242 (3.3)	2/256 (0.8)
2	$1.8 \times 10^3 (0.18)$	$1.8x10^3 (0.18) 1.9x10^3 (0.19)$	7/127 (5.5)	6/123 (4.8)
4	$1.6 \times 10^3 (0.16)$	$1.6x10^3 (0.16) 3.3x10^3 (0.33)$	(0.9) 66/9	8/105 (8.5)
000	1.9x10 ³ (0.19)	$1.9 \times 10^3 (0.19) 3.6 \times 10^3 (0.36)$	7/56 (12.5)	12/60 (13.1)

PCT/EP99/03075

15

20

25

Example 7

Muscle regeneration by genetically-modified fibroblasts expressing MyoD

Fibroblasts isolated from human fetal skin were expanded and transduced in vitro with a high-titer stock of the retroviral vector LBSN, carrying a lacZ gene encoding a cytoplasmic form of β-galactosidase (Salvatori, G., et al., 1993).

Transduction efficiency, estimated by the number of β -gal-staining cells in culture, ranged between 40 and 70%, making further selection of the transduced cells (e.g., G418 resistance) unnecessary. Transduced fibroblasts were exposed to the adeno-MyoD vector at an m.o.i. of 2,000, and then injected into the regenerating TA muscles of scid/bg mice (106 cells/muscle in a single injection).

The efficiency of myogenic conversion of transduced fibroblasts was checked by allowing part of the cell culture to differentiate in vitro. On average, about 70% of the fibroblasts underwent myogenic conversion in these conditions, and most myotubes (> 90 %) scored positive for cytoplasmic \(\mathbb{B}\)-gal expression. After 1, 2 and 4 weeks, mice were sacrificed and the TA serially sectioned and tibialis anterior muscles were cut and stained for \(\mathbb{B}\)-gal activity.

Two weeks after injection, \(\beta - gal + \) fibers were observed in 7 out of 8 injected muscles. Higher magnification clearly revealed fibers accumulating the reporter gene product at variable levels, suggesting a variability in the proportion of injected/host cells in the \(\beta - gal + \) fibers.

The presence and contribution of human cells in the newly formed fibers was confirmed by immunocytochemical analysis using an antibody which recognizes human but not mouse fetal myosin heavy chains, which

A North Spirits on

13

10

15

20

25

WO 99/58646 PCT/EP99/03075

showed patches of intensive expression of human muscle-specific proteins in the regenerating areas.

In general, the average number of B-gal+ fibers per muscle obtained by injecting human converted fibroblasts was lower than that 5 - observed after injection of murine fibroblasts, although comparable to the number of positive cells observed in experiments in which lacZ-transduced human satellite cells were used. This indicates that human cells are, as expected, less efficient than murine ones in colonizing a mouse muscle.

However, and more importantly, human converted fibroblasts perform like primary myogenic cells in vivo, in terms of both number and size of newly formed fibers.

In order to better quantify the survival of human myogenically converted fibroblasts in mouse muscle, the Applicant also performed dot-blot Southern analysis with an Alu probe, as previously described (Salvatori, G., et al., 1993). To this purpose DNA was extracted from TA muscles of scid/bg mice two weeks after the injection of MyoD-converted human fibroblasts (following the same experimental protocol described above). As a positive control, the same number of human satellite cells was injected in similarly treated contra-lateral TA muscles.

The results of the dot-blot indicated that approximately 0.1% of the DNA extracted from TA injected with converted fibroblasts was of human origin; on the other hand this value raised to 0.3% in the case of the sample injected with human satellite cells.

Because histochemical analysis had revealed that virtually all the ß-galactosidase staining was inside muscle fibers, we can conclude that the observed values faithfully reflect the percentage of human nuclei

25

incorporated into regenerated muscle fibers.

Example 8

Immune response against cells exposed ex vivo to the adenoviral vector.

PCT/EP99/03075

One of the major problems associated to the in vivo use of adenoviral vectors in immunocompetent recipients is the induction of a significant, T-cell-mediated immune response against both the viral proteins and the product of the transgene(s). In the model described herein, no viral particle is injected directly in vivo. However, injected cells had been exposed shortly before administration to the adenoviral vector in vitro.

Replication-defective adenoviral DNA is expected to be rapidly diluted, and eventually lost, in cells in active cell division. Expression of the MyoD transgene, however, blocks almost immediately cell division in the converted fibroblasts, thus preventing the possibility of diluting the adenoviral vector by continuous culture.

To test whether an immune response could be raised by cells expressing a defective adenoviral genome in vivo, the Applicant injected 2×10^6 C3H murine skin fibroblasts converted to myogenesis by exposure to the adeno-MyoD vector in the regenerating muscle of singeneic mice.

Cells were labelled with DiI before injection and not with a lacZ transgene to avoid a possible immune reaction against the \(\beta \)-galactosidase protein. Animals were sacrificed 7, 14 and 21 days after the injection, and muscles analysed for the presence of immune infiltrate by immunofluorescence using antibodies against cell surface markers of murine leukocytes and macrophages.

No significant immune infiltrate was detected for up to 3 weeks after injection around the labelled cells, even though by this time all treated mice had developed antibodies against Ad5 proteins, which recognise adeno-infected but not control C3H fibroblasts

These experiments indicate that fibroblasts converted in vitro by exposure to an adeno-MyoD vector, and administered in vivo by intramuscular injection, do not elicit, and are therefore unlikely to be eliminated by, a cell-mediated cytotoxic immune response.

5

i ja In contrast, direct injection of Adenoviral vector into regenerating muscle induced a strong immune infiltrate, as already reported (Yang, Y., et al., 1996).

15

25

BIBLIOGRAPHY

Salvatori, G., G. Ferrari, A. Mezzogiorno, S. Serivdei, M. Coletta, P. Tonali, R. Giavazzi, G. Cossu, and F. Mavilio. 1993. Retroviral vector-mediated gene transfer into human primary myogenic cells leads to expression in muscle fibers in vivo. Human Gene Ther. 4:713-723.

Webster, C. e, H.M. Blau 1990. Accelerated age-related decline in replicative life span of Duchenne muscular dystrophy myoblasts: implication for cell and gene therapy. Somatic Cell Mol. Genet. 16:557-565.

Davis, R. L., H. Weintraub, and A.B. Lassar. 1987. Expression of a single transfected cDNA converts fibroblasts to myoblasts. Cell 51:987-1000.

Cossu, G., S. Tajbakhsh, and M. Buckingham. 1996. Myogenic specification in mammals. Trends Genet. 12:218-223.

Cossu, G. 1997. Unorthodox myogenesis: possible developmental significance and implications for tissue histogenesis and regeneration. Histol. Histopathol. 12:755-760.

Breton, M., Z., Li, D. Paulin, J.A. Harris, F. Rieger, M. Pincon-Raymond, and L. Garcia. 1995. Myotube driven myogenic recruitment of cells during in vitro myogenesis. Develop. Dynam. 202:126-136.

Gibson, A.J., J. Karasinski, J. Relvas, J. Moss, T.G. Sherratt, P.N. Strong, and D.J. Watt. 1995. Dermal fibroblasts convert to a myogenic lineage in mdx mouse muscle. J. Cell Science 108:207-214.

Salvatori, G., L. Lattanzi, M. Coletta, S. Aguanno, E. Vivarelli, R. Kelly, G. Ferrari, J. A. Harris, F. Mavilio, M. Molinaro, and G. Cossu. 1995. Myogenic conversion of mammalian fibroblasts induced by differentiating muscle cells. J. Cell Science 108:2733-2739.

10

15

20

Murry, C. E., M. A. Kay, T. Bartosek, S. D. Hauschka, and S. M. Schwartz. 1996. Muscle differentiation during repair of myocardinal necrosis in rats via gene transfer with MyoD. J. Clin. Invest. 98:2209-2217.

PCT/EP99/03075

Cusella - De Angelis, M.G., S. Molinari, A. Ledonne, M. Coletta, E. Vivarelli, M. Bouchè, M. Molinaro, S. Ferrari, and G. Cossu, 1994. Differential response of embryonic and fetal myoblasts to TGFß: a possible regulatory mechanism of skeletal muscle histogenesis. Development 120:925-933.

Bader, D., T. Masaki, and D. A. Fischman. 1982. Immunochemical analysis of myosin heavy chain during avian myogenesis in vivo and in vitro. J. Cell Biol. 95:763-770.

Hasty, P., A. Bradley, J. H. Morris, D. G. Edmondson, J. M. Venuti, E. N. Olson, and W. H. Klein. 1993. Muscle deficiency and neonatal death in mice with a targeted mutation in the myogenin gene. Nature 375:787-790.

Edom, F., V. Mouly, J. P. Babert, M. Y. Fiszman, and G. S. Butler-Browne. 1994. Clones of human satellite cells can express in vitro both fast and slow myosin heavy chains. Develop. Biology 164:219-229.

Yasin, R., K. C. Van Beers, C. E. Nurse, S. Al-Ani, D. N. Landon, and E. J. Thompson. 1977. A quantitative technique for growing human adult skeletal muscle in culture starting from mononucleated cells. J. Neurol. Sci. 32:347-360.

Chomczynski, P., and N. Sacchi. 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chlorophorm extraction.

Anal. Biochem. 162:156-159.

Bober, E., G.E. Lyons, T. Braun, G. Cossu, M. Buckingham, and

the first of the feet of the feet of the first of the fir

5

10

20

H.H. Arnold. 1991. The muscle regulatory gene Myf-6 has a biphasic pattern of expression during early mouse development. J. Cell Biol. 113:1255-1265.

Boulter, J., W. Luyten, K. Evans, P. Mason, M. Ballivet, D. Goldman, D. Stengelin, S. Martin, S. Heinemann, and J. Patric, 1985. Isolation of a clone coding for the α-subunit of a mouse acetylcholine receptor. J. Neurosci. 5:2545-2552.

Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual. 2nd ed., Cold Spring Harbor Laboratory Press, New York.

Yang, Y., S. E. Haecker, Q. Su, and J. Wilson, 1996. Immunology of gene therapy with adenoviral vectors. Hum. Mol. Genet. 5:1703-1712.

Tajbakhrsh, S., E. Vivarelli, M. G. Cusella-De Angelis, D. Rocancourt, M. Buckingham, and G. Cossu 1994. A population of myogenic cells derived from the mouse neural tube. Neuron 13:813-821.

Lyons, G. E., S. Muhlebach, A. Moser, R., Masood, B. M. Paterson, M. Buckingham, and J. C. Perriard. 1991. Development regulation of creatin kinase gene expression by myogenic factors in embryonic mouse and chick skeletal muscle. Development 113:1017-1029.

CLAIMS

- 1. A method for the preparation of genetically-modified fibroblasts expressing a muscle lineage commitment gene, which comprises:
- a) ex-vivo transduction of fibroblasts with a therapeutic gene or a gene capable of correcting a gene defect;
 - b) transient expression of the muscle lineage commitment gene in fibroblasts transduced as at point (a), through transformation of the cells with a high-efficiency DNA transfer method, wherein the muscle lineage commitment gene is under the control of a strong promoter.
 - A method according to claim 1, wherein the therapeutical gene is the dystrophin gene.
 - A method according to claim 1, wherein the high-efficiency DNA transfer method is a viral vector.
 - A method according to claim 3, wherein said viral vector is selected from baculovirus, adeno-related viruses, adeno-virus.
 - 5. A method according to claim 3, wherein said vector is an adenovirus.
 - A method according to claim 1, wherein the muscle lineage commitment gene is selected from MyoD, Myf-5, MRF4 and myogenin.
- 20 7. A method according to claim 6, wherein said gene is MyoD.
 - A method according to claim 1, wherein the muscle lineage commitment gene is under the control of a viral promoter.
 - Genetically-modified fibroblasts obtainable by the method of claims 1-
- 25 10. Fibroblasts according to claim 9, wherein the muscle lineage commitment gene is MyoD.

Nixon & Vanderhye P.C. (12/97)

RULE 63 (37 C.F.R. 1.63) DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

I am the subject n	original, first and sole ir natter which is claimed a	reby declare that my residence, nventor (if only one name is list and for which a patent is sought	ed below) or ar	original, first and joint invent	tated below no or (if plural na	ext to my name, and I believe ames are listed below) of the
		ts and the use thereof				
	Ification of which (check attached hereto	applicable box(s)):				
	attached nereto as filed on		!! 0 4	lication Serial No.		
	as filed as PCT Internati	and application No.	as 0.5. App.			
		application) was amended on	FCI/EF99	/03075 on <u>05</u>	-05-1999	
(ii cip	philodolic to o.g. of 1 o 1	application) was unchase on				
amendm with 37 (listed bel which pri Priority F	ent referred to above. C.F.R. 1.56. I hereby of low and have also identi- ority is claimed or, if no- foreign Application(s):	ewed and understand the cont I acknowledge the duty to disc claim foreign priority benefits ur fied below any foreign application priority is claimed, before the fil	lose information nder 35 U.S.C. on for patent or ing date of this	n which is material to the pat 119/365 of any foreign appli- inventor's certificate having a application:	entability of to cation(s) for p filing date be	his application in accordance vatent or inventor's certificate fore that of the application or
MI98A00	ion Number		Countr	y		onth/Year Filed
WII96AUU	11004		Italy			08-05-1998
I hereby Applicat	claim the benefit under a ion Number	35 U.S.C. §119(e) of any United	States provision Date/Month/Ye		<i>i</i> .	
the subje	ct matter of each of the 12, I acknowledge the o	35 U.S.C. 120/365 of all prior claims of this application is not duty to disclose material informed international filing date of the control	disclosed in su ation as define	ch prior applications in the ma	anner provide	d by the first paragraph of 35
	S./PCT Application(s): ion Serial No.		Day/Month/Ye	ar Filed		Status: patented pending, abandoned
Section 10 And on be riumber (owner's/or patent: A 22770; M 33363; Jet Alan M. K 41090; M afforney n	not of Title 18 of the Unitehalf of the owner(s) here to the owner where 3 attorneys to prosecuthur R. Crawford, 2532; and E. Nusbaum, 32348; Joffry H. Nelson, 30481; Joffry H. Nelson, 30431; John Cagen, 36178; Robert A. Nichelle N. Lester, 32331; antes/numbers no longer	ade with the knowledge that willife ded States Code and that such will of, I hereby appoint NIXON & V. all communications are to be discussed in the supplication and to transact_Larry S. Nixon_25599; Robert_Gichael I. Keenan_32106_Bryan Fin R. Lastova_3120_Bryan Fin R. Lastova_3120_Bryan Fin folian_2834_B. J. Sarforf_36653. Frank P. Presst_1928_Joseph Switch the firm and to act and rely s Nixon & Vandethye on behalf of the supplies of the sup	ful false stateme: ANDERHYE P. ANDERHO, and the et all business in A. Vanderhye, I. Davidson, 302 Burnam, Jr. 2936 James D. Berqu I. Presta, 35329; olely on instruct	ats may jeopardize the validity of C., 1100 North Glebe Rd., 8th following attorneys thereof (of the Patent and Trademark Offs 27076; James T. Hosmer, 3018- 51; Stanley C. Spooner, 27393; 6; Thomas E. Byrne, 32205; Maist, 34776; Updeep S. Gill, 3733. Joseph A. Rhon, 37515. Lake.	of the application of the application of the same addresses connected to the connected of the control of the co	on or any patent issued thereon on M. A 2394–1414, telephone is) individually and collectively herewith and with the resulting aris, 31552; Richard G. Besha chard, 2900%; Duane M. Byers 299554–1, Scott Davidson, 31489, hea. 34275; Donald L. Jackson on & Vanderhye to delete any ssignee, attorney, firm, or other
1.	Inventor's Signature: Inventor:	Fulvio	7		_ Date: _1	3-10-2000
. \	inventor.	(first)	MI	MAVILIO		Italian
1/21	Residence: (city)	Milano (IIISI)	IVII	(last) (state/country) Italy		(citizenship)
10	Post Office Address:	Via Olgettina, 58 – Milano, It	alv	(state/codility) Italy		
~	(Zip Code)	20132	ui)			
	(Lip code)	20102				
2.	Inventor's Signature:				Date:	
	Inventor:				—	1
		(first)	MI	(last)		(citizenship)
	Residence: (city)			(state/country)		(,
	Post Office Address:					
	(Zip Code)					
3.	Inventor's Signature:				Date:	
		(first)	MI	(last)		(citizenship)
	Residence: (city)	()	****	(state/country)		(Guzensnip)
	Post Office Address:					
	(Zip Code)					

FOR ADDITIONAL INVENTORS, check box 🔲 and attach sheet with same information and signature and date for each.